

DISCRETE POLYLOGARITHM FUNCTIONS

TOM CUCHTA^{1,2} — DALLAS FREEMAN²

¹Dept. of Mathematics, Marshall University, Huntington, West Virginia, USA

²Dept. of Computer Science and Mathematics, Fairmont State University, Fairmont, West Virginia, USA

ABSTRACT. We investigate a discrete analogue of the polylogarithm function. Difference and summation relations are obtained, as well as its connection to the discrete hypergeometric series.

1. Introduction

Discrete special functions and their applications have been the topic of numerous papers in recent years, see e.g., discrete analogues of Bessel and hypergeometric functions [2, 3, 6], analogues of orthogonal polynomials [4, 5, 8], semidiscrete multivariable models [12–14], and discrete models of physics [1].

On the other hand, the classical polylogarithm function

$$\mathcal{L}i_s(t) = \sum_{k=1}^{\infty} \frac{t^k}{k^s}.$$

has seen numerous applications in diverse areas such as Fermi-Dirac integrals [16], conformal field theory [11], thermoelectrics [17], and blackbody radiation [15].

We are interested in expanding the theory of discrete special functions to include discrete polylogarithm functions, which we define by

$$\text{Li}_s(t; n, \xi) = \sum_{k=1}^{\infty} \frac{t^{nk} \xi^k}{k^s}, \quad (1)$$

where $s, \xi \in \mathbb{C}$ and $n \in \mathbb{R}$, which is the discrete analogue of the polylogarithm function.

© 2023 Mathematical Institute, Slovak Academy of Sciences.

2020 Mathematics Subject Classification: 33E30, 33C99.

Keywords: discrete special functions, polylogarithm, hypergeometric series.

This research was made possible by NASA West Virginia Space Grant Consortium, Training Grant #NNX15AI01H.



Licensed under the Creative Commons BY-NC-ND 4.0 International Public License.

2. Preliminaries and definitions

The forward difference operator Δ is defined by $\Delta f(t) = f(t + 1) - f(t)$. The discrete shift for $n, m \in \{0, 1, \dots\}$ is the relation

$$t^n(t - n)^m = t^{n+m}. \quad (2)$$

The discrete fundamental theorem of calculus is

$$\sum_{k=a}^b \Delta f(k) = f(b + 1) - f(a). \quad (3)$$

The classical generalized hypergeometric series is defined by

$${}_p\mathcal{F}_q(\mathbf{a}; \mathbf{b}; t) = \sum_{k=0}^{\infty} \frac{(a_1)_k \dots (a_p)_k}{(b_1)_k \dots (b_q)_k} \frac{t^k}{k!},$$

where $\mathbf{a} = (a_1, \dots, a_p)$ and $\mathbf{b} = (b_1, \dots, b_q)$ for some constants $a_i, b_j \in \mathbb{C}$. The discrete hypergeometric series is

$${}_pF_q(\mathbf{a}; \mathbf{b}; t, n, \xi) = \sum_{k=0}^{\infty} \frac{(a_1)_k \dots (a_p)_k}{(b_1)_k \dots (b_q)_k} \frac{t^{nk} \xi^k}{k!}.$$

It is known [3, Proposition 2] that the ${}_pF_q$ and ${}_p\mathcal{F}_q$ are related by

$${}_pF_q(\mathbf{a}; \mathbf{b}; t, n, \xi) = {}_{p+n}\mathcal{F}_q(\mathbf{a}, \mathbf{t}; \mathbf{b}; \xi(-n)^n), \quad (4)$$

where

$$\mathbf{t} = \left(\frac{-t}{n}, \frac{-t+1}{n}, \dots, \frac{-t+n-1}{n} \right).$$

The falling factorials are defined in terms of the Γ function by

$$a^{\underline{b}} = \frac{\Gamma(a+1)}{\Gamma(a-b+1)}, \quad (5)$$

and ratios of Γ functions obey the following known asymptotic relationship [7]

$$\frac{\Gamma(x+\beta)}{\Gamma(x)} \sim x^\beta \quad (6)$$

The related Pochhammer symbols are defined for $a \in \mathbb{C}$ by

$$(a)_k = a(a+1) \dots (a+k-1).$$

The polylogarithm obeys many interesting formulas that can be found in the books [9, 10]. We now express many such properties for the $\mathcal{L}i$:

$$\mathcal{L}i_n(t) = t_{n+1}\mathcal{F}_n(1, 1, \dots, 1; 2, 2, \dots, 2; t), \quad (7)$$

$$t \frac{\partial \mathcal{L}i}{\partial t} = \mathcal{L}i_{s-1}(t), \quad (8)$$

$$\mathcal{L}i(t) = \int_0^t \frac{\mathcal{L}i_s(\tau)}{\tau} d\tau, \quad (9)$$

and

$$\sum_{j=0}^{p-1} \mathcal{L}i_s \left(z e^{\frac{2\pi i j}{p}} \right) = p^{1-s} \mathcal{L}i_s(z^p). \quad (10)$$

3. Discrete polylogarithms

If $t \in \{0, 1, \dots\}$, then the series (1) converges, since the factor t^{nk} will ultimately vanish for sufficiently large k . Now we establish convergence for complex t .

THEOREM 3.1. *If $t \notin \{0, 1, \dots\}$ and $n \in \{0, 1, 2, \dots\}$, then the series (1) converges for $|t| < \sqrt[n]{\frac{1}{|\xi|}}$.*

Proof. We assume that $t \notin \{0, 1, 2, \dots\}$ because such t -values cause the series to terminate due to the factor of t^{nk} in the summand. To apply the ratio test, set $a_k = \frac{t^{nk} \xi^k}{k^s}$ and consider the limit

$$\begin{aligned} \lim_{k \rightarrow \infty} \left| \frac{a_{k+1}}{a_k} \right| &= \lim_{k \rightarrow \infty} \left| \frac{t^{nk+n} \xi^{k+1} k^s}{(k+1)^s t^{nk} \xi^k} \right| \\ &= \lim_{k \rightarrow \infty} \left| \frac{\Gamma(t - nk + 1) \xi k^s}{\Gamma(t - nk - n + 1) (k+1)^s} \right| \\ &\approx \lim_{k \rightarrow \infty} \left| \xi \left(\frac{k}{k+1} \right)^s t^{-nk+1 - (-nk-n+1)} \right| \\ &= \lim_{k \rightarrow \infty} \left| \xi \left(\frac{k}{k+1} \right)^s t^n \right| = |\xi t^n|. \end{aligned}$$

Hence the series converges whenever $|t| < \sqrt[n]{\frac{1}{|\xi|}}$, completing the proof. \square

When the subscript is a non-negative integer, the series (1) reduces to a discrete hypergeometric function, analogous to (7).

THEOREM 3.2. *If $m, n \in \{0, 1, \dots\}$, then*

$$\text{Li}_m(t; n, \xi) = \xi t^n {}_{m+1}F_m(1, 1, \dots, 1; 2, 2, \dots, 2; t - n, n, \xi). \quad (11)$$

Proof. Using (2), compute

$$\begin{aligned} \xi t^{\underline{n}} F_m(1, \dots, 1; 2, \dots, 2; t-n, n, \xi) &= \xi t^{\underline{n}} \sum_{k=0}^{\infty} \frac{(1)_k \dots (1)_k (t-n)^{\underline{nk}} \xi^k}{(2)_k \dots (2)_k k!} \\ &= \sum_{k=0}^{\infty} \frac{(k!)^{m+1}}{((k+1)!)^m} \frac{t^{n(k+1)} \xi^{k+1}}{k!} = \sum_{k=1}^{\infty} \frac{t^{nk} \xi^k}{k^m} = \text{Li}_m(t; n, \xi), \end{aligned}$$

completing the proof. \square

The previous theorem implies a representation as a classical generalized hypergeometric series via (4).

COROLLARY 3.3. *If $m, n \in \{0, 1, \dots\}$, then*

$$\text{Li}_m(t; n, \xi) = \xi t^{\underline{n}} {}_{m+1+n}\mathcal{F}_m(1, 1, \dots, 1, \mathbf{t}; 2, \dots, 2; \xi(-n)^{\underline{n}}),$$

where $\mathbf{t} \in \mathbb{R}^{1 \times n}$ with

$$\mathbf{t} = \left(\frac{-t+n}{n}, \frac{-t+n+1}{n}, \dots, \frac{-t+2n-1}{n} \right).$$

The following theorem is a discrete analogue of (8).

THEOREM 3.4. *The functions (1) obey the formula*

$$t \Delta \text{Li}_s(t-1; n, \xi) = n \text{Li}_{s-1}(t; n, \xi). \quad (12)$$

Proof. Compute

$$\begin{aligned} t \Delta \text{Li}_s(t-1; n, \xi) &= t \Delta \sum_{k=1}^{\infty} \frac{\xi^k (t-1)^{\underline{nk}}}{k^s} = tn \sum_{k=1}^{\infty} \frac{\xi^k (t-1)^{\underline{nk-1}}}{k^{s-1}} \\ &= n \sum_{k=1}^{\infty} \frac{\xi^k t^{\underline{nk}}}{k^{s-1}} = n \text{Li}_{s-1}(t; n, \xi), \end{aligned}$$

completing the proof. \square

The following corollary is a discrete analogue of (9).

COROLLARY 3.5. *The functions (1) obey the formula*

$$\text{Li}_s(t-1; n, \xi) = n \sum_{k=1}^{t-1} \frac{\text{Li}_{s-1}(k; n, \xi)}{k}.$$

Proof. Divide (12) by t and sum to obtain

$$\sum_{k=1}^{t-1} \Delta \text{Li}_s(k-1; n, \xi) = n \sum_{k=1}^{t-1} \frac{\text{Li}_{s-1}(k; n, \xi)}{k}.$$

Since $\text{Li}_s(0; n, \xi) = 0$, applying (3) on the left-hand side yields

$$\text{Li}_s(t-1; n, \xi) = n \int_1^t \frac{\text{Li}_{s-1}(\tau; n, \xi)}{\tau} \Delta\tau.$$

Recognizing the integral on the right-hand side as a sum completes the proof. \square

The summation (10) has the following discrete analogue.

THEOREM 3.6. *If $p \in \{1, 2, 3, \dots\}$, then*

$$\sum_{j=0}^{p-1} \text{Li}_n \left(t, n, \xi e^{\frac{2\pi ij}{p}} \right) = p^{1-n} \text{Li}_n(t, np, \xi^p).$$

Proof. Calculate

$$\sum_{j=0}^{p-1} \text{Li}_s \left(t, n, \xi e^{\frac{2\pi ij}{p}} \right) = \sum_{j=0}^{p-1} \sum_{k=1}^{\infty} \frac{t^{nk} \xi^k e^{\frac{2\pi k ij}{p}}}{k^s} = \sum_{k=1}^{\infty} \frac{t^{nk} \xi^k}{k^s} \sum_{j=0}^{p-1} e^{\frac{2\pi k ij}{p}}$$

By the well-known sum of roots of unity

$$\sum_{j=0}^{p-1} e^{\frac{2\pi k ij}{p}} = \begin{cases} p, & k|p, \\ 0, & k \nmid p. \end{cases}$$

we obtain

$$\begin{aligned} \sum_{j=0}^{p-1} \text{Li}_s \left(t, n, \xi e^{\frac{2\pi ij}{p}} \right) &= p \sum_{k=1, k|p}^{\infty} \frac{t^{nk} \xi^k}{k^s} = p \sum_{\ell=1}^{\infty} \frac{t^{n p \ell} \xi^{p \ell}}{(p \ell)^s} \\ &= p^{1-s} \sum_{\ell=1}^{\infty} \frac{t^{(np)\ell} (\xi^p)^\ell}{\ell^s} = p^{1-s} \text{Li}_s(t, np, \xi^p), \end{aligned}$$

completing the proof. \square

4. Conclusion

We have established discrete analogues of many of the properties of the polylogarithm functions. Future work can expand into looking at applications of these functions, understanding special cases such as representations when n is a negative integer, and other discrete analogues such as inverse tangent integrals and the Legendre χ function.

Acknowledgement. This research was made possible by NASA West Virginia Space Grant Consortium, Training Grant #NNX15AI01H.

REFERENCES

- [1] BAS, E.—OZARSLAN, R.—YILMAZER, R.: *Spectral structure and solution of fractional hydrogen atom difference equations*, AIMS Mathematics **5** (2020), 1359–1371.
- [2] BOHNER, M.—CUCHTA, T.: *The Bessel difference equation*, Proc. Am. Math. Soc. **145** (2017), 1567–1580.
- [3] BOHNER, M.—CUCHTA, T.: *The generalized hypergeometric difference equation*, Demonstr. Math. **51** (2018), 62–75.
- [4] CUCHTA, T.—PAVELITES, M.—TINNEY, R.: *The Chebyshev difference equation*, Mathematics **8** (2020) no. 1, DOI:10.3390/math8010074.
- [5] CUCHTA, T.—LUKETIC, R.: *Discrete hypergeometric Legendre polynomials*, Mathematics **9**, (2021), no. 20, DOI: 103390/math9202546.
- [6] CUCHTA, T.—GROW, D.—WINTZ, N.: *Discrete matrix hypergeometric functions*, J. Math. Anal. Appl. **518**(2) (2023), no 2, <https://doi.org/10.1016/j.jmaa.2022.126716>
- [7] ERDÉLYI, A.—TRICOMI, F.: *The asymptotic expansion of a ratio of gamma functions*, Pacific J. Math **1** (1951), 133–142.
- [8] GANIE, J.—JAIN, R.: *Basic analogue of Legendre polynomial and its difference equation*, Asian J. Math. Stat. **12** (2019), 1–7.
- [9] LEWIN, L.: *Polylogarithms and Associated Functions*. (With a foreword by A. J. Van der Poorten). North-Holland Publishing Co., New York-Amsterdam, 1981.
- [10] LEWIN, L.: Editor. *Structural properties of polylogarithms*. Mathematical Surveys and Monographs, Vol. 37. American Mathematical Society, Providence, RI, (1991).
- [11] SACHDEV, S.: *Polylogarithm identities in a conformal field theory in three dimensions*, Phys. Lett. B **309** (1993), 285–288.
- [12] SLAVÍK, A.: *Discrete Bessel functions and partial difference equations*, J. Difference Equ. Appl. **24** (2018), 425–437.
- [13] SLAVÍK, A.: *Asymptotic behavior of solutions to the semidiscrete diffusion equation*, Appl. Math. Lett. **106** (2020), <https://doi.org/10.1016/j.aml.2020.106392>
- [14] SLAVÍK, A.: *Spatial maxima, unimodality, and asymptotic behaviour of solutions to discrete diffusion-type equations*, J. Difference Equ. Appl. **28** (2022), 126–140.
- [15] STEWART, S.: *Blackbody radiation functions and polylogarithms*, J. Quant. Spectrosc. Radiat. Transf. **113** (2012), 232–238.
- [16] ULRICH, M.—SENG, W.—BARNES, P.: *Solutions to the Fermi-Dirac integrals in semiconductor physics using Polylogarithms*, J. Comput. Electronics **1** (2002), 431–434.
- [17] MOLLI, M.—VENKATARAMANIAN, K.—VALLURI, S.: *The polylogarithm and the Lambert W functions in thermoelectrics*, Canadian J. Physics **89** (2011), 1171–1178.

Received September 30, 2022

Department of Mathematics
Marshall University
1 John Marshall Drive
Huntington, West Virginia 25755
USA
E-mail: cughta@marshall.edu

Dept. of Computer Science and Mathematics
Fairmont State University
Fairmont, West Virginia 26554
USA
E-mail: dallas.freeman@gmail.com